



congenital bile acid synthesis defect type 2

Congenital bile acid synthesis defect type 2 is a disorder characterized by cholestasis, a condition that impairs the production and release of a digestive fluid called bile from liver cells. Bile is used during digestion to absorb fats and fat-soluble vitamins, such as vitamins A, D, E, and K. People with congenital bile acid synthesis defect type 2 cannot produce (synthesize) bile acids, which are a component of bile that stimulate bile flow and help it absorb fats and fat-soluble vitamins. As a result, an abnormal form of bile is produced.

The signs and symptoms of congenital bile acid synthesis defect type 2 often develop in infancy. Affected infants usually have a failure to gain weight and grow at the expected rate (failure to thrive) and yellowing of the skin and eyes (jaundice) due to impaired bile flow and a buildup of partially formed bile. Excess fat in the feces (steatorrhea) is another feature of congenital bile acid synthesis defect type 2. As the condition progresses, affected individuals can develop liver abnormalities including inflammation or chronic liver disease (cirrhosis). Some individuals with congenital bile acid synthesis defect type 2 cannot absorb certain fat-soluble vitamins, which can result in softening and weakening of the bones (rickets) or problems with blood clotting that lead to prolonged bleeding.

If left untreated, congenital bile acid synthesis defect type 2 typically leads to cirrhosis and death in childhood.

Frequency

The prevalence of congenital bile acid synthesis defect type 2 is unknown. Together, all congenital defects of bile acid synthesis are thought to have a prevalence of 1 to 9 per million people.

Genetic Changes

Mutations in the *AKR1D1* gene cause congenital bile acid synthesis defect type 2. The *AKR1D1* gene provides instructions for making an enzyme called 3-oxo-5- β -steroid 4-dehydrogenase. This enzyme is found in certain liver cells that produce bile acids. Bile acids are produced from cholesterol in a multi-step process. The 3-oxo-5- β -steroid 4-dehydrogenase enzyme is responsible for the third step in that process, which converts 7 α (α)-hydroxy-4-cholesten-3-one to 7 α -hydroxy-5 β -cholesten-3-one.

AKR1D1 gene mutations result in a 3-oxo-5- β -steroid 4-dehydrogenase enzyme with severely reduced function. Without enough functional enzyme, the conversion

of 7 α -hydroxy-4-cholesten-3-one to 7 α -hydroxy-5 β -cholesten-3-one is impaired. The 7 α -hydroxy-4-cholesten-3-one instead gets converted into abnormal bile acid compounds that cannot be transported out of the liver into the intestine, where the bile acids are needed to absorb fats and fat-soluble vitamins. As a result, cholesterol and abnormal bile acids build up in the liver and fat-soluble vitamins are not absorbed, which contribute to the signs and symptoms of congenital bile acid synthesis defect type 2.

Inheritance Pattern

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- CBAS2
- cholestasis with delta(4)-3-oxosteroid 5-beta-reductase deficiency

Diagnosis & Management

Genetic Testing

- Genetic Testing Registry: Bile acid synthesis defect, congenital, 2
<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1856127/>

Other Diagnosis and Management Resources

- Cleveland Clinic: Cirrhosis of the Liver
<http://my.clevelandclinic.org/health/articles/cirrhosis-of-the-liver>
- MedlinePlus Encyclopedia: Cholestasis
<https://medlineplus.gov/ency/article/000215.htm>

General Information from MedlinePlus

- Diagnostic Tests
<https://medlineplus.gov/diagnostictests.html>
- Drug Therapy
<https://medlineplus.gov/drugtherapy.html>
- Genetic Counseling
<https://medlineplus.gov/geneticcounseling.html>

- Palliative Care
<https://medlineplus.gov/palliativecare.html>
- Surgery and Rehabilitation
<https://medlineplus.gov/surgeryandrehabilitation.html>

Additional Information & Resources

MedlinePlus

- Encyclopedia: Bile
<https://medlineplus.gov/ency/article/002237.htm>
- Encyclopedia: Cholestasis
<https://medlineplus.gov/ency/article/000215.htm>
- Health Topic: Jaundice
<https://medlineplus.gov/jaundice.html>
- Health Topic: Liver Diseases
<https://medlineplus.gov/liverdiseases.html>

Genetic and Rare Diseases Information Center

- Congenital bile acid synthesis defect, type 2
<https://rarediseases.info.nih.gov/diseases/10045/congenital-bile-acid-synthesis-defect-type-2>

Additional NIH Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: Cirrhosis
<https://www.niddk.nih.gov/health-information/liver-disease/cirrhosis>

Educational Resources

- Disease InfoSearch: Bile Acid Synthesis Defect, Congenital, 2
<http://www.diseaseinfosearch.org/Bile+Acid+Synthesis+Defect%2C+Congenital%2C+2/831>
- Merck Manual Home Edition: Cholestasis
<http://www.merckmanuals.com/home/liver-and-gallbladder-disorders/manifestations-of-liver-disease/cholestasis>
- Merck Manual Home Edition: Cirrhosis of the Liver
<http://www.merckmanuals.com/home/liver-and-gallbladder-disorders/fibrosis-and-cirrhosis-of-the-liver/cirrhosis-of-the-liver>

- Orphanet: Congenital bile acid synthesis defect type 2
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=79303
- Orphanet: Disorder of bile acid synthesis
http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=79168

Patient Support and Advocacy Resources

- American Liver Foundation
<http://www.liverfoundation.org/>
- CORE: Fighting Gut and Liver Disease (UK)
<http://corecharity.org.uk/>
- Council for Bile Acid Deficiency Diseases
<http://www.bileacid.org/>
- National Organization for Rare Disorders (NORD)
<https://rarediseases.org/rare-diseases/bile-acid-synthesis-disorders/>
- Resource List from the University of Kansas Medical Center: Liver Conditions
<http://www.kumc.edu/gec/support/liver.html>

ClinicalTrials.gov

- ClinicalTrials.gov
<https://clinicaltrials.gov/ct2/results?term=%22congenital+bile+acid+synthesis+defect+type+2%22+%5BDISEASE%5D+OR+%22cholestasis+with+delta%28%29-3-oxosteroid+5-beta-reductase+deficiency%22+%5BDISEASE%5D+OR+%22inborn+error+of+bile+acid+synthesis%22+%5BDISEASE%5D+OR+NCT00571272+%5BID-NUMBER%5D+OR+NCT01589523+%5BID-NUMBER%5D>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28bile+acid+synthesis+defect%29+AND+%28AKR1D1%29+OR+%28delta4-3-oxosteroid+5beta-reductase+deficiency%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>

OMIM

- BILE ACID SYNTHESIS DEFECT, CONGENITAL, 2
<http://omim.org/entry/235555>

Sources for This Summary

- Clayton PT. Disorders of bile acid synthesis. J Inherit Metab Dis. 2011 Jun;34(3):593-604. doi: 10.1007/s10545-010-9259-3. Epub 2011 Jan 13. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21229319>
 - Lemonde HA, Custard EJ, Bouquet J, Duran M, Overmars H, Scambler PJ, Clayton PT. Mutations in SRD5B1 (AKR1D1), the gene encoding delta(4)-3-oxosteroid 5beta-reductase, in hepatitis and liver failure in infancy. Gut. 2003 Oct;52(10):1494-9.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12970144>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1773813/>
 - Mindnich R, Drury JE, Penning TM. The effect of disease associated point mutations on 5 β -reductase (AKR1D1) enzyme function. Chem Biol Interact. 2011 May 30;191(1-3):250-4. doi: 10.1016/j.cbi.2010.12.020. Epub 2010 Dec 24.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21185810>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3101292/>
-

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/condition/congenital-bile-acid-synthesis-defect-type-2>

Reviewed: April 2015

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services